

**ENGLISH TRANSLATION**

**Only For Use in Animal Health.**

**1 NAME OF THE BIOLOGICAL VETERINARY MEDICINAL PRODUCT**

ALPHA DIP 2000 ,Formalin inactivated dip suspension against Vibriosis and Pasteurellosis in seabass

**2 PHARMACEUTICAL FORM**

Dip suspension

**3 STATEMENT AND POWER OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)**

Formalin inactivated bacteria cultures of:

<i>Listonella anguillarum</i> (serotype O1)	RPS ≥ 75
<i>Photobacterium damsela</i> subsp. <i>piscicida</i>	RPS ≥ 60

**4 INDICATIONS**

Prevents Clinical symptoms and mortality caused by *Listonella anguillarum* serotype O1 (Vibriosis) and *Photobacterium damsela* subsp. *piscicida* (pasteurellosis). The product is intended for the first and second vaccination.

**5 CONTRAINDICATIONS**

None

**6 ADVERSE EFFECTS**

Due to handling stress, vaccination may be followed by a temporary reduced appetite subsequent to vaccination.

**7 TARGET SPECIES**

Sea bass

**8 DOSAGE FOR EACH SPECIES**

Dilute one litre of vaccine with 9 litres of clean rearing water. Immerse the fish into the diluted vaccine suspension for 30 seconds in the knotless net. Feeding should be withheld at least one day before the vaccination. Temperate the vaccine to the room temperature and shake the vaccine bottle prior to use.

**9 METHOD AND ROUTE OF ADMINISTRATION**

The following vaccination scheme is recommended:

First and second vaccination is applied by dipping (ALPHA DIP 2000) of fish with average weight approximately 1 gram and 5 grams respectively.

Primary vaccination is intended for fish residing in the hatchery that average 1 g in weight. Vaccinate minimum 2 weeks prior to transfer to sea cages. Temperate the vaccine to room temperature and shake the vaccine bottle prior to use.

Dilute one litre of vaccine with 9 litres of clean rearing water (1:10 dilution). Immerse the fish into the diluted vaccine for 30 seconds. Discard the vaccine when a total of 20 kg fish have been immersed per 10 litre of the diluted vaccine.

Second Vaccination is intended for fish residing in sea cages that average 5 g in weight.

Temperate the vaccine and shake the vaccine bottle prior to use.

Dilute one litre of vaccine with 9 litres of clean rearing water (1:10 dilution). Immerse the fish into the diluted vaccine for 30 seconds. Discard the vaccine when a total of 100 kg fish have been immersed per 10 litre of the diluted vaccine.

## **10 LEGAL WITHDRAWAL PERIOD**

**Zero days**

## **11 COMMERCIAL PACKAGING INFORMATION:**

1000 ml, White HDPE bottles, in carton boxes of 10 units

## **12 SPECIAL STORAGE PRECAUTIONS AND SHELF LIFE**

Store in dark at 2-8 °C. Do not freeze. Protect from sunlight.  
Use the entire content when first opened within the same working day.  
Do not use after the expiry date stated on the label.  
Shelf life is 24 months.

## **13 SPECIAL PRECAUTIONS**

In any population there will be a small number of individuals, which fail to respond well to vaccination. Occasional mortality may occur if individuals fail to respond or the immune system is suppressed by concurrent infections, poor nutritional status, genetic factors or other stressful environmental conditions.

## **14 ADVICE ON CORRECT ADMINISTRATION FOR VACCINATORS AND VETERINARIES**

Vaccinate only fish free of clinical symptoms of disease, demonstrating normal behavior. For fish safety, avoid vaccinating at water temperature higher than 25°C and during periods with rapidly increasing or decreasing water temperatures.

Shake well the vaccine bottle prior to use.

**15 SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS DERIVED FROM SUCH VETERINARY MEDICINAL PRODUCTS**

Any unused veterinary medicinal product or waste material derived from such products should be disposed with help of disinfectants or by burning.

**16 WARNINGS**

Keep out of the reach and sight of children .

Consult to a veterinarian if an unexpected effect is observed.

Please refer to the packaging insert for detailed information about the use of the vaccine before use

**17 NAME AND ADDRESS OF THE LICENCE HOLDER**

Manufacturer  
PHARMAQ AS  
Industrivegen 50  
N-7863 Overhalla  
NORWAY

Importer company  
PHARMAQ Veteriner Eczacı Deposu ve Su Ürünleri Ticaret Limited Şirketi.  
Address: KARACAOĞLAN MAH.6166 SK. 21/A BORNOVA  
Izmir 35070, Türkiye  
Tel. 232 422 23 10  
Fax 232 422 32 27

**18 MARKETING AUTHORISATION DATE AND NUMBER**

11.05.2015 / P-0076

**19 DATE ON WHICH THE PACKAGE INSERT WAS LAST REVISED**

July 2015

**20 OTHER INFORMATION**

For any further information about this veterinary medicinal product, please contact your local license holder company officials.